

QUALI-PRO

Fosetyl-Al 80 WDG

Fungicide

ACTIVE INGREDIENT: % **BY WT.**
 Aluminum tris (O-ethyl phosphonate): . . . 80.0%
INERT INGREDIENTS: 20.0%
TOTAL: 100.0%

EPA Reg. No. 66222-161 EPA Est. No. 37429-GA-001^{BT}
 37429-GA-002^{EO}

Letter(s) in lot number correspond(s)
 to superscript in EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:
Makhteshim Agan of North America, Inc.
 4515 Falls of Neuse Road, Suite 300
 Raleigh, NC 27609

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call Prosur at 1-877-250-9291.	

QUALI-PRO

Fosetyl-Al 80 WDG

Fungicide

ACTIVE INGREDIENT: % BY WT.
Aluminum tris (O-ethyl phosphonate): . . . 80.0%
INERT INGREDIENTS: 20.0%
TOTAL: 100.0%

EPA Reg. No. 66222-161 EPA Est. No. 37429-GA-001^{BT}
37429-GA-002^{BD}

Letter(s) in lot number correspond(s)
to superscript in EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:
Makhteshim Agan of North America, Inc.
4515 Falls of Neuse Road, Suite 300
Raleigh, NC 27609

13375

Net Contents: 5.5 Pounds

EPA 010908/
Notif 011309/Rev C

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call Prostar at 1-877-250-9291.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult agency responsible for pesticide regulation.

This product is not to be used on home or residential sites.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted-entry interval of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of the treated areas until sprays have dried.

Quali-Pro Fosetyl-AI 80 WDG is a systemic fungicide that, when used according to the label directions and as part of good management practices programs, will control or suppress listed fungal diseases in a variety of crops.

COMPATIBILITY

Turf and Ornamentals: Some foliar fertilizers and copper-based fungicides are not compatible with Quali-Pro Fosetyl-AI 80 WDG, and a tank mix of these products with Quali-Pro Fosetyl-AI 80 WDG may result in phytotoxicity to various plants. To avoid this phytotoxicity, apply Quali-Pro Fosetyl-AI 80 WDG at least 14 days before or after an application of a copper-based fungicide; and in these spray programs, avoid applying with air blast equipment. Stickers, extenders, and wetting agents must not be mixed with Quali-Pro Fosetyl-AI 80 WDG. Do not tank mix Quali-Pro Fosetyl-AI 80 WDG with the following products because they are not compatible with Quali-Pro Fosetyl-AI 80 WDG: Daconil 2787 Flowable Fungicide (or other registered flowable formulations containing chlorothalonil), Fore Fungicide flowable formulations (or other registered flowable formulations containing mancozeb), Bayleton® 50W (or other registered similar formulations containing

triadimefon), Rubigan™ A.S. (or other registered flowable formulations containing fenamirol), Spectro™ 90WDG (or other similar registered formulations containing thiophanate-methyl and chlorothalonil).

Physical Compatibility Test: Before mixing Quali-Pro Fosetyl-AI 80 WDG with any other product including but not limited to soluble concentrate or flowable formulations, conduct a physical compatibility test. Use a clear container (1 pint) and mix a small amount of spray solution by adding each component in the order and ratio as required for making large amounts of the tank mix solution. Stir the contents and allow to settle for 5-15 minutes. Solutions that stay in suspension or can be remixed readily are considered compatible. Increased compatibility may result if a buffering agent is used.

TOLERANCE

Although Quali-Pro Fosetyl-AI 80 WDG has been found to be safe to use with the species listed on this label, it is impossible to test all species or varieties under all types of growing conditions for tolerance to Quali-Pro Fosetyl-AI 80 WDG. Always test for plant phytotoxicity on a small scale using the recommended rates before preparing spray solutions or applying to large areas of plants.

PREPARATION OF SPRAY SOLUTIONS

Only prepare the amount of spray solution required for one day's application.

1. Always use clean water when preparing spray tank solutions. Ensure the spray tank is clean prior to use. Add clean water to fill the tank 1/4 to 1/2 full.
2. With the agitator running, slowly add the required amount of Quali-Pro Fosetyl-AI 80 WDG to the tank.
3. Slowly add an alkaline buffering agent when required (see **TANK MIXES** below).
4. Fill the spray tank with the remaining amount of water and continue agitation until the application is completed.

TANK MIXES

Plant phytotoxicity from solubilized copper has been observed under several situations:

- 1) when products containing copper are tank-mixed with Quali-Pro Fosetyl-AI 80 WDG, **OR**
- 2) when unbuffered Quali-Pro Fosetyl-AI 80 WDG solutions are applied to plant foliage that bear copper residues from prior treatments with copper products. Do not tank mix Quali-Pro Fosetyl-AI 80 WDG with copper compounds.

In situations when copper compounds are applied **prior to** or will be applied **after** applications of Quali-Pro Fosetyl-AI 80 WDG, increase the pH of the Quali-Pro Fosetyl-AI 80 WDG tank solution (see Step 3 under **PREPARATION OF SPRAY SOLUTIONS** above) to 6.0 or above using an alkaline buffer. Examples of alkaline buffers include potassium carbonate (use 3 lbs. for every 5 lbs. Quali-Pro Fosetyl-AI 80 WDG) and diammonium phosphate (use 5 lbs. for every 5 lbs. Quali-Pro Fosetyl-AI 80 WDG). Use caution when using potassium carbonate as the buffer because it reacts during the addition to form carbon dioxide gas (CO₂). Do not seal the spray tank until all of the CO₂ gas is released, a process that usually takes about 5 minutes.

APPLICATION DIRECTIONS FOR FOLIAR SPRAYS

For systemic control of various fungal diseases, apply Quali-Pro Fosetyl-AI 80 WDG according to these label directions.

Apply Quali-Pro Fosetyl-AI 80 WDG in a sufficient amount of water to ensure thorough coverage of plant foliage. The spray volumes will vary by crop and growth stage.

APPLICATION DIRECTIONS FOR SPRINKLER IRRIGATION SYSTEMS

Do not apply Quali-Pro Fosetyl-AI 80 WDG through chemigation in California.

Apply this product only through sprinkler irrigation systems including mini-sprinkler or micro-jet, drip, solid set, and center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: A suspension of Quali-Pro Fosetyl-AI 80 WDG is prepared in a mixing tank by adding 1/2 to 3/4 of the desired amount of water. Start the agitator and then add the required amount of Quali-Pro Fosetyl-AI 80 WDG followed by the remaining volume of water. Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of Quali-Pro Fosetyl-AI 80 WDG into the irrigation water line to deliver the desired rate per acre. The suspension should be injected with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

NOTE: For mini-sprinkler and drip irrigation systems, avoid field irrigation over the treated area for 24 to 48 hours after completion of the Quali-Pro Fosetyl-AI 80 WDG treatment.

For solid set and center pivot irrigation systems, after completion of the Quali-Pro Fosetyl-AI 80 WDG treatment, do not irrigate the treated area until the foliage has dried to prevent wash-off of the chemical from the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

- Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.
- The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled and drained.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

ORNAMENTALS AND BEDDING PLANTS

Quali-Pro Fosetyl-AI 80 WDG controls diseases caused by:

- Pythium and Phytophthora in ornamentals and bedding plants grown in field nursery, greenhouse, landscaping and conifer nursery situations, and
- downy mildew and fire blight in ornamentals.

Quali-Pro Fosetyl-AI 80 WDG suppresses bacterial blight caused by certain pathogens of *Xanthomonas campestris*.

Time applications to occur before signs of disease appear. Use Quali-Pro Fosetyl-AI 80 WDG along with good cultural management practices. To avoid plant injury, follow these label directions and do not apply Quali-Pro Fosetyl-AI 80 WDG above the recommended rates or at shorter spray intervals than recommended in this label.

ORNAMENTALS

Application Method:	Foliar
Typical Ornamentals:	Aglaonema, Aphelandra, Azalea, Bougainvillea, Boxwood, <i>Cattelya skinneri</i> , Cissus, Dieffenbachia, Hibiscus, Juniper, Leather-leaf Fern, Pittosporum, Philodendron, Pothos, Rhododendron, Spathiphyllum, Taxus media
Rate:	2.5 – 5.0 lbs. product per 100 gals. water
Specific Directions:	Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Use a rate of 2.5 – 5.0 lbs. product/A when applying in spray volumes of less than 100 gals./A. Do not exceed 400 gals. of spray solution per acre. Make repeat applications as needed but not more often than every 30 days.
Application Method:	Drench
Typical Ornamentals:	Aphelandra, Azalea, Boxwood, Cissus, Dieffenbachia, Japanese Holly, Juniper, Monterey Pine, Philodendron, Pieris, Pittosporum, Rhododendron, Schefflera, Spathiphyllum, Taxus media
Rate:	0.4 – 0.8 lbs. product (6.4 – 12.8 oz.) per 100 gals. water
Specific Directions:	Apply spray solution to 400 sq. ft. (2 pts. solution per sq. ft.) of plants. Make repeat applications as needed but not more often than every 30 days.
Application Method:	Soil incorporation – to control Phytophthora species only
Typical Ornamentals:	Azalea, Pieris, or Rhododendron
Rate:	0.5 – 0.8 lbs. product (8 – 12.8 oz.) per cubic yard of soil
Specific Directions:	Mix product with soil immediately before potting. Soil incorporate only to plants with an established root system. Foliar or drench applications may be made when disease development is expected, but do not apply more often than every 30 days.

BEDDING PLANTS

Application Method:	Foliar
Typical Ornamentals:	Begonia, Pansy, Vinca, Marigold, Zinnia, Petunia, Geranium, Impatiens
Rate:	1.25 – 4.0 lbs. product per 100 gals. water
Specific Directions:	Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Do not exceed 400 gals. of spray solution per acre. Make repeat applications as needed but not more often than every 14 days.

TANK MIXES FOR ORNAMENTAL AND BEDDING PLANTS

Quali-Pro Fosetyl-AI 80 WDG is compatible with most commonly used fungicides. Quali-Pro Fosetyl-AI 80 WDG will provide control of additional diseases (such as Aerial Web Blight, Alternaria Leaf Blight, Botrytis Blight, Fusarium Leaf Spot, Helminthosporium Leaf Spot, and Ray Blight) when tank mixed with other fungicides such as CHIPCO® 26019 fungicide or other iprodione-containing fungicides. Consult the label of the tank mix product for use directions, the diseases controlled, and application rates. When another fungicide is used in combination with Quali-Pro Fosetyl-AI 80 WDG, follow all label directions for the use of the other product. Make applications at the recommended rate for control of the target disease organism.

Refer to the section above on **COMPATIBILITY** for a list of products that cannot be tank mixed with Quali-Pro Fosetyl-AI 80 WDG.

CONIFER NURSERIES*

Disease:	Phytophthora Root Rot
Application Method:	Dip
Typical Conifers:	Douglas firs, spruce, and pines
Rate:	2.5 lbs. product per 100 gals. water
Specific Directions:	Thoroughly wet plant and roots by dipping into the Quali-Pro Fosetyl-AI 80 WDG solution immediately prior to transplanting.

NOTE: During the dipping process, users must wear chemical/water-resistant gloves, goggles or face shield, long pants (coveralls), long-sleeved shirt, and shoes plus socks.

Disease:	Phytophthora Root Rot
Application Method:	Foliar
Typical Ornamentals:	Douglas firs, spruce, and pines
Rate:	2.5 – 5.0 lbs. product per 100 gals. water
Specific Directions:	Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Use a rate of 2.5 – 5.0 lbs./A when applying in spray volumes of less than 100 gals./A. Do not exceed 400 gals. of spray solution per acre. Make repeat applications as needed but not more often than every 30 days.

*Not registered for use in California.

ROSES

Disease:	Downy Mildew (<i>Peronospora sparsa</i>)
Application Method:	Foliar – field, container, landscape, miniature varieties
Rate:	2.5 lbs. product per 100 gals. water
Specific Directions:	Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Use a rate of 2.5 – 5.0 lbs./A when applying in spray volumes of less than 100 gals./A. Do not exceed 400 gals. of spray solution per acre. Make repeat applications as needed but not more often than every 7 days.
NOTE:	Use Quali-Pro Fosetyl-AI 80 WDG in combination with a sanitation program designed to prevent the spread of this disease to healthy plants.

FIRE BLIGHT – SUPPRESSION ONLY

Typical Ornamentals:	Ornamental pear, pyracantha, hawthorn
Application Method:	Foliar
Rate:	2.5 lbs. product per 100 gals. water
Specific Directions:	Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Use a rate of 2.5 – 5.0 lbs./A when applying in spray volumes of less than 100 gals./A. Do not exceed 400 gals. of spray solution per acre. Applications must be made early (start at pre-bloom) to prevent Fire Blight infection. Make repeat applications as needed until blooming ends but not more often than every 7 days.
NOTE:	Use Quali-Pro Fosetyl-AI 80 WDG in combination with a sanitation program designed to prevent the spread of this disease to healthy plants.

BACTERIAL BLIGHT – SUPPRESSION ONLY*

- Disease:** *Xanthomonas campestris* pathovars *hederae*, *dieffenbachiae*, *syngonii*, and *fici* – Suppression Only
- Typical Ornamentals:** English ivy, schefflera, anthurium, dieffenbachia, spathiphyllum, syngonium, and ficus
- Application Method:** Foliar
- Rate:** 2 – 4 lbs. product per 100 gals. water
- Specific Directions:** Apply Quali-Pro Fosetyl-AI 80 WDG to wet the foliage. Use a rate of 2.5 – 5.0 lbs./A when applying in spray volumes of less than 100 gals./A. Do not exceed 400 gals. of spray solution per acre. Make repeat applications as needed but not more often than every 7 days.

NOTE: Use Quali-Pro Fosetyl-AI 80 WDG in combination with a sanitation program designed to prevent the spread of this disease to healthy plants. Read the section on **COMPATIBILITY** and the use of Quali-Pro Fosetyl-AI 80 WDG with copper fungicides or other compounds.

*Not registered for use in California.

TURF – QUALI-PRO FOSETYL-AL 80 WDG ALONE

- Diseases:** Pythium diseases including blight and root rot, Yellow Tuft
- Application Method:** Foliar
- Typical Sites:** Turf grass on golf courses, sod farms, commercial turf areas
Not for use on home or residential turf
- Rate:** 4.0 oz. product per 1000 sq. ft., 14-day intervals
- Rate:** 8.0 oz. product per 1000 sq. ft., 21-day intervals
- Specific Directions:** Apply Quali-Pro Fosetyl-AI 80 WDG in 1 to 5 gals. of water per 1000 sq. ft. Agitate solutions during the spray operations. Calibrate equipment before making application. Use in a season-long program to control listed diseases by applying when conditions favor disease. Make repeat applications at the spray interval listed based on use rate. Allow the foliage to dry before watering or mowing treated areas.

RESTRICTIONS:

- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock and poultry.

TURF – QUALI-PRO FOSETYL-AL 80 WDG AND TANK MIX PARTNERS

Application Method: Foliar

Typical Sites: Turf grasses on golf courses, sod farms, industrial and municipal turf areas

Not for use on home or residential turf

Spray Intervals: 14 days

Diseases: Summer Stress Complex/Summer Decline (*Pythium*, *Anthracnose*, *Rhizoctonia spp.*) AND/OR Anthracnose (*Colletotrichum graminicola*)

Tank Mix Products:

QP Fosetyl AI 80 WDG

plus

26GT

or

Daconil Ultrex®

or

Manicure™ DG

or

Dithane® T/O

or

Protect T/O

or

Mancozeb DG

or

Fore® WP

or

Heritage®

or

Banner® MAXX®

or

ProStar® 70WP

Diseases:

Rate (oz. product per 1000 sq. ft.)

4.0 – 8.0

plus

3.0 – 4.0

or

3.2

or

3.3

or

8.0

or

8.0

or

8.0

or

4.0 – 8.0

or

0.2 – 0.4

or

1.0 – 2.0

or

1.5 – 3.0

Bentgrass Deadspot (*Ophlosphaerella agrostis*)

Tank Mix Products:

Quali-Pro Fosetyl-AI 80 WDG

plus

Fore® Rainshield™

or

Daconil Ultrex®

Rate (oz. product per 1000 sq. ft.)

4.0

plus

8.0

or

3.2

(continued on next page)

(continued from previous page)

Specific Directions: Use the most restrictive of label limitations for Quali-Pro Fosetyl-AI 80 WDG or tank mix partner. Always apply at the recommended rates. Do not mix with products that prohibit such mixing. Apply in 1 to 5 gals. of water per 1000 sq. ft. Agitate solutions during the spray operations. Calibrate equipment prior to use. Repeat at recommended intervals. Allow foliage in treated areas to dry before watering or mowing. To improve the control of anthracnose, use Quali-Pro Fosetyl-AI 80 WDG in a preventative, multi-application program in which applications begin prior to outbreak of the disease. To improve the control of deadspot, use Quali-Pro Fosetyl-AI 80 WDG in a preventative, multi-application program in which applications begin prior to outbreak of the disease and when conditions favor development of the disease.

NOTE: When used in turf management programs, Quali-Pro Fosetyl-AI 80 WDG promotes turf that is thick and vigorous and tolerant to such stress factors as heat and humidity. This healthy grass is less susceptible to diseases and other turf pests such as weeds and algae.

RESTRICTIONS:

- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock and poultry.

The following fungicides are known to be compatible with Quali-Pro Fosetyl-AI 80 WDG and any one may be used in place of products listed above for use as a tank mix with Quali-Pro Fosetyl-AI 80 WDG to control Summer Stress Complex:

Bayleton® 50*	Fungo® 50 WSB
Cavalier 50WSB	MANhandle™
Cavalier Flowable	Manicure™ 6 Flowable
Chlorostar	Manicure™ T/O Flowable
Cleary's 3336™ WP*	Manicure™ Ultrex®
Cleary's 3336™ F*	Medallion®
Compass™	Pentathlon®
Concorde™ DF	Pentathlon® DF
Concorde™ SST	Revere™ 4000
Consyst	SysStar®
Curalan® EG	SysTec® 1998
Daconil® Weather Stik®*	SysTec® 1998 WDG
Daconil ZN® Rainshield NT™	Sythane® WSP
Eagle®	Terrachlor® 75WP and 400*
Echo® 90DF	Terraguard® 50W
Echo® Flowable	Terrazole 35% WP
Endorse™	Touche EG
Fluid Fungicide	Turfcide®400
Fungo® Flo	TwoSome™ Flowable Fungicide

***NOTE:** When preparing spray tank mixtures, add these products to the spray tank and allow complete dispersion in water before adding Quali-Pro Fosetyl-AI 80 WDG.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container and keep tightly closed. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable Container (flexible-bag-all weights): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid-fifty lbs or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid-greater than fifty lbs): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product

container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

CHIPCO, Bayleton, Compass, and Prostar are registered trademarks of the Bayer Group. Endorse™ is a trademark of Arvesta.

Banner Maxx, Heritage, Daconil, Daconil Ultrex, Weather Stik, Daconil ZN, and Medallion are registered trademarks of Syngenta.

Dithane, Systhane, Eagle and Fore are registered trademarks of Dow AgroSciences Company.

Cavalier, Manicure, MANhandle, Revere, TwoSome and Lesco are trademarks of Lesco, Inc. Curalan is a registered trademark of BASF Corporation.

3336, Spectro and Protect are trademarks of Cleary Chemical Corp.

Consyst, Systec and Syster are registered trademarks of Regal Chemical Company.

Echo is a registered trademark of SIPCAM AGRO USA, Inc.

Fungo is a registered trademark of the Scotts Company.

Terrachlor, Terraguard, Terrazole and Turfcide are registered trademarks of Uniroyal Chemical Company, Inc.

Concorde is a trademark and Pentathlon is a registered trademark of the DuPont Company.

Quali-Pro® is a registered trademark of Makhteshim Agan of North America, Inc.

Rubigan A.S. is a registered trademark of Gowan Company, LLC.

Eagle Fungicide is a registered trademark of Dow AgroSciences, LLC.

