

FUNGICIDE

FLUAZINAM 40SC SELECT™

BROAD-SPECTRUM TURF FUNGICIDE.



Fluazinam 40SC Select™ is a broad-spectrum, multi-site fungicide to control diseases in turf grasses on golf courses only. The active ingredient, fluazinam, is the only registered fungicide for turf in the FRAC 29 group and, as a multi-site contact fungicide, carries a very low risk of developing resistance. **Fluazinam 4SC Select** is a great rotational option to chlorothalonil based fungicides for season-long protection against the toughest turf diseases.

PRODUCT FEATURES

- Unique multi-site contact active ingredient.
- Excellent addition to resistance-management programs.
- Excellent control of dollar spot and additional turf diseases.
- Perfect tank-mix partner with systemics for inside out protection.
- Controls dollar spot that is resistant to other chemistries.

LABELED USES

- Golf course tees, greens, fairways, and rough.

LABELED DISEASES

Algal scum	Gray snow mold	Pink patch
Anthracnose	Large patch	Pink snow mold
Brown blight	Leaf spot	Red thread
Brown patch	Melting out	Stem rust
Dollar spot		

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

40% fluazinam

FRAC GROUP

29

FORMULATION

Suspension concentrate

CHEMICAL FAMILY

2,6-dinitroanilines

COMPARE TO

Secure®

MODE OF ACTION

Oxidative phosphorylation
uncoupler

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

4 x 0.5 gal, 2 x 2.5 gal

EPA REG NUMBER

89442-37

FLUAZINAM 40SC SELECT™

BROAD-SPECTRUM TURF FUNGICIDE.

APPLICATION INFORMATION

Apply **Fluazinam 40SC Select™** in sufficient water to obtain adequate, uniform coverage of the turf.

- Apply at 0.5 fl. oz. per 1,000 sq. ft. (6 acres per gallon) every 14 days.
- Apply after mowing with sufficient water to uniformly cover the turf and allow area to dry completely before irrigation.
- Refer to the product label for additional information about this product and tank mixing.
- Not for use on home lawns or any residential turf.

Turfgrass disease control

- Apply **Fluazinam 40SC Select** in a preventative disease control program.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed area to completely dry before mowing.
- For control of foliar diseases, allow sprayed area to completely dry before irrigation.
- For optimum turf quality and disease control, use **Fluazinam 40SC Select** in conjunction with turf management practices that promote good plant health.
- Evaluate spray additives prior to use. Label directions are based on data obtained with no additives.
- Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

Restrictions

- **DO NOT** exceed 253 fl oz of **Fluazinam 40SC Select** (8.4 pounds active ingredient) per acre, per year.
- **DO NOT** apply more than twelve (12) applications at 21.5 fl oz per acre (0.5 fl oz/1000 sq ft), per year. For best results, use spray mixture the same day it is prepared.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Fluazinam 40SC Select is a trademark of Albaugh, LLC. Secure is a registered trademark of Ishihara Sangyo Kaisha. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.